

Evaluation of Bioactive compounds and antimicrobial analysis of the essential oil from the leaves of *Sauropus androgynous* in Rajasthan India

Alagbe, Olujimi John *

Department of Animal Nutrition and Biochemistry, Sumitra Research Institute, Gujarat, India.

Corresponding author: Olujimi John, Department of Animal Nutrition and Biochemistry, Sumitra Research Institute, Gujarat, India.

Received date: November 06, 2025; **Accepted date:** November 17, 2025; **Published date:** November 29, 2025

Citation: Alagbe, Olujimi John, (2025), Evaluation of Bioactive compounds and antimicrobial analysis of the essential oil from the leaves of *Sauropus androgynous* in Rajasthan India, *J. Nutrition and Food Processing*, 8(10); DOI:10.31579/2637-8914/340

Copyright: © 2025, Olujimi John. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract:

Sauropus androgynous essential oil contains unique bioactive compounds with therapeutic potentials and it delivers powerful antioxidant, anti-inflammatory, antidiabetic, cytotoxic, gastro-protective, immune-protective, cardio-protective, antimicrobial, antidiarrheal, anti-helminthic amongst others. GC-MS analysis of *Sauropus androgynous* essential oil revealed the presence of 33 bioactive compounds were recorded representing 96.27 % composition dominated by Squalene (19.63 %), Caryophyllene (13.92 %), Nonane, 4-methyl (13.82 %) and n-hexadecanoic acid (10.45 %). Other bioactive compounds examined had a concentration less than 3 %. A synergy between the major and minor compounds are working together to protect, heal, restore and improve the overall vitality of the body. The result obtained on the antimicrobial activity of *Sauropus androgynous* essential oil revealed a high percentage inhibition 77.81 %, 72.00 %, 80.09 %, 73.73 % and 76.26 % against *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Shigella flexineri* and *Salmonella typhi* respectively. The results compare well with that of the standard drug (Ciprofloxacin) (65.00 -71.00 %). It was concluded that *Sauropus androgynous* essential oil contains measurable concentration of compounds that can disrupt the cell membrane of pathogens and will also help to address the increasing cases of antimicrobial resistance.

Key words: bioactive; compounds; antimicrobials; free radicals; oxidative stress

Introduction

Sauropus androgynous, an evergreen, multipurpose herb belongs to the Phyllanthaceae plant family which is found in the tropical rain forests of West Africa, although it is also widely distributed in parts of Asia, South America and Australia (Purba and Paengkoum, 2012; Bao et al., 2020). Inside its leaves are bioactive compounds that go far beyond nutrition (Juxian and Xian, 2022; Hernandez and Alagbe, 2025a). The leaves are rich in minerals like calcium, magnesium and iron supporting bone strength and electrolyte balance (Singh et al., 2011; Singh et al., 2021), vitamins (B-complex and C) and protein between 6 – 10 % (Lin et al., 1999; Basker, 2012). For centuries, Traditional healers have used its leaves to soothe infection, skin disorder, pyrexia, bronchitis, diarrhea and relief from acute and chronic infections because its tissues are filled with therapeutic potentials (Samad et al., 2013; John, 2024b).

The plant contains a unique profile of flavonoids and phenolic acids, nature's compounds known to neutralize the activities of free radicals, reinforces the cardiovascular system by strengthening blood vessel, acts as liver protectors, metabolic regulators, reducing inflammation and restoring enzyme balance (Alagbe, 2024; Singh et al., 2022). They also contain alkaloids, tannins, steroids and saponins which possess several pharmacological functions which includes, cytotoxic, antimicrobial, antitumor, antidiabetic, anti-cancer, antifungal, immune-modulatory,

gastro-protective, derma-protective, anti-helminthic, antidiarrheal, anti-allergic, antihistaminic, antipyretic, antiseptic, antispasmodic, antiulcer and antiviral (Naemsuvan, 2013; Zhang et al., 2022). Methanolic extract from *Sauropus androgynous* leaves and stems show high radical scavenging free activity (Gonzalez et al., 2022; Purba et al., 2022), measurable compounds inhibit the activities of bacteria cells in microbial assays (Caldeira et al., 2004; Chatrou et al., 2012). Its poultices can be used for skins and wounds (Gao et al., 2014; Musa et al., 2020). Aqueous extract suppresses inflammatory cytokines, reducing airway infections and regulating sugar absorption in the body (Adewale et al., 2021; John, 2024a).

Previous studies by Rajeswari et al. (2018) have shown that *Sauropus androgynous* leaf extract contained 9,12,15-Octadecatrienoic acid at 12.90 %, Ethyl 9,12,15-Octadecatrienoate (15.93 %), Ethyl (9Z,12Z)-9,12-Octadecadienoate (11.28 %), Hexadecanoic acid (13.76 %), 1-(+)-Ascorbic acid 2,6-Dihexadecanoate (27.81 %) as the prominent bioactive compounds during GC-MS analysis. Senthamarai and Anusha (2012) also reported that GC-MS profiling in the leaves of *Sauropus Androgynus* showed that it contained 2(1H) Naphthalenone (41.17 %), Azulene (36.20 %), Pyrene, hexadecahydro (9.07 %), squalene (8.06 %) and 1, 14-Tetradecanediol (2.82 %) as major bioactive compounds. Previous report

has shown that for every newly developed antibiotic, bacterial strains rapidly develop resistance mechanisms against it. Therefore, there is an urgent need for alternative ways to treat bacterial infections and to combat the problem of antimicrobial resistance (Carranza et al. 2015; Hernandez and Alagbe, 2025b). One of the alternatives that can be utilised for combating antimicrobial resistance is traditional herbal remedies from plants. Evaluating the bioactive compounds in *Sauropus androgynous* oil will further help to unveil their pharmacological and therapeutic properties.

Materials and methods

Location of the experiment, plant collection and extraction of *Sauropus androgynous* oil

This study was carried out at the Microbiology Department Gandhi College of Agriculture, Rajasthan India between October to December 2025. The Institution is situated between 21°35'N 26°09'E East India. *Sauropus androgynous* fresh leaves were harvested from Rajasthan India and sent to the department of botany, Gandhi College of Agriculture for identification before it was registered under voucher number HU09/2025C/008. The identified leaves were sorted and shade dried for 13 days until a constant weight was achieved. Dried *Sauropus androgynous* leaves were pulverized using mechanical grinder. Extraction of oil was done by hydrodistillation with a Clevenger-type apparatus according to the procedures outlined by Singh et al. (2021). 250 g of the pulverized *Sauropus androgynous* was added to 1000 mL of water heated in a glass flask at 60 °C for 20 minutes, steam passes via the condenser and when cooled it was collected in a beaker. The oil collected by decantation at the end of the distillation was filtered, dried over column of anhydrous sodium sulfate, and introduced into glass bottles and stored in a refrigerator at 4 °C.

Gas chromatography – mass spectrometry analysis of *Sauropus androgynous* essential oil

GC/MS analysis of *Sauropus androgynous* essential oil was done using Claudius 5006 GC-MS Auto Sampler (China) equipped with two silica capillary columns, interfaced with a quadrupole detector (single quadrupole acquisition Method-MS parameters report), source temperature 230°C, Quadrupole temperature 150 °C; the temperature program was 60 °C for 2 min, 60-240 °C at 3 °C/min, then kept at 240 °C during 8 min; injector temperature, 240 °C. The mass spectrometry transfer line temperature, 250 °C; carrier gas, helium at a flow rate of 0.7 ml/min; injection type, split, 20:1; ionization voltage, 70 eV; electron multiplier 1000 eV; scan range 33-400 amu; scan rate, 1.56 scan/s.

Identification of components Interpretation on mass spectrum GC-MS was conducted using the database of National Institute Standard and Technology (NIST, 2001) having more than 62,000 patterns. The spectrum of the unknown component was compared with the spectrum of the known components stored in the NIST library.

Antimicrobial assay

The antimicrobial activity of essential oil of *Sauropus androgynous* essential oil was screened against five standard strains of stock bacteria from the Microbiology department, Gandhi College of Agriculture, Rajasthan. *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Shigella flexineri* and *Salmonella typhi*. Microplate Alamar Blue Assay was used to determine susceptibility or resistance of the essential oils to all the selected bacteria strains. Organisms were grown in Mueller Hinton broth and inoculums were adjusted to 0.5 McFarland standard. Essential oil (20 µg/mL) was added in the wells; control wells do not contain essential oil. The volume of 96-well plate was made up to 200 µL. Finally, 5 x 10⁶ cells were added in all wells including both control and test. The plate was sealed with parafilm and incubated for 18 - 20 hours. Alamar Blue Dye was dispensed in each well and shaken at about 80 revolution per minute in a shaking incubator for 2 – 3 hours.

Plates were covered with foil in shaking incubator. Change in color of Alamar Blue dye from blue to pink indicated the growth in bacterial strains. Ciprofloxacin was used as the reference drug.

Bioactive compounds in *Sauropus androgynous* essential oil by GC-MS analysis is presented in Table 1. A total of 33 bioactive compounds were recorded representing 96.27 % composition dominated by Squalene (19.63 %), Caryophyllene (13.92 %), Nonane, 4-methyl (13.82 %) and n-hexadecanoic acid (10.45 %). These compounds have been previously associated with anti-inflammatory (Sinha and Munshi, 2011; Namkeleja et al., 2014), antioxidant (Omokore and Alagbe, 2019; Oluwafemi et al., 2021), cytotoxic and immuno-stimulatory properties (Alagbe, 2022; Muritala et al., 2022). According to Shittu and Alagbe (2020), Squalene has the capability to neutralize free radicals' activities in the body, support blood vessels integrity and protects the heart, lungs and nervous systems. Caryophyllene have been reported to be a proven powerhouse, kills pathogens and inhibits fungal growth (Tiwari et al., 2011; Ojediran et al., 2024). Hexadecanoic acid excels in lowering blood sugar and supports liver detoxification (Sharma, 2012; Jing-Chung et al., 2007; Lima et al., 2010). Nonane, 4-methyl can interfere with mitochondrion energy production in cancer cells, fights infection and calm the nervous system (Choi et al., 2013). All the 33 bioactive compounds have numerous pharmacological properties making them useful in traditional medicine for the treatment of acute and chronic inflammation, cough, fever, gastrointestinal disorders, skin infection, eye problems, sexually transmitted infections, stomach ulcers, pain, snake bite, tooth ache amongst others (Alagbe et al., 2022; Agubosi et al., 2022). A previous study by Senthamarai and Anusha (2012) on the GC-MS analysis of *Sauropus androgynous* leaf reveals that squalene and phytol contained a concentration of 8.06 % and 0.88 % respectively. Lee et al. (2011) also recorded a lower concentration of 0.12 % and 3.07 % for phytol and hexadecanoic acid. Rajeswari et al. (2018); Basker (2012) reported that *Sauropus androgynous* leaf extract contained a higher concentration of 13.80 % hexadecanoic acid. Samad et al. (2018) reported 10.40 % and 0.04 % for hexadecanoic acid and 9, 17-Octadecadienol, and these values were almost similar to the outcome in this experiment. However, geographical location, age of plant, specie, storage method, harvesting technique, processing methods have been identified as factors that influence the concentration of bioactive compounds in medicinal plants (Awa et al., 2012). A synergy between the minor and major components in essential oil from *Sauropus androgynous* in this study forms a rich network of phytochemicals that reduces inflammation by suppressing pro-inflammatory cytokines (Momza et al., 2012; Alagbe et al., 2022), produces bronchodilator effect by opening respiratory airway thereby making breathing easier (Zubairu et al., 2025), improve cardio-vascular health and electrolyte balance thus improving the overall health and vitality of the body (Alagbe, 2025).

The result obtained on the antimicrobial activity of *Sauropus androgynous* essential oil revealed a high percentage inhibition 77.81 %, 72.00 %, 80.09 %, 73.73 % and 76.26 % against *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Shigella flexineri* and *Salmonella typhi* respectively. The results compare well with that of the standard drug (Ciprofloxacin) (65.00 -71.00 %). The presence of squalene, the prominent bioactive compounds in the oil might have been responsible for their antimicrobial activity. Squalene contains measurable compounds that inhibits the activities of some bacteria's (Ajiboye et al., 2013). Phyto-components like hexadecanoic acid, caryophyllene, cyclopentaneundecanoic acid, 8-octadecenoic acid, humulene, gamma- Terpinene-3-Carene, 9, 17-Octadecadienol, Undecane, 2,10-dimethyl, 2-methyl-Glutaric acid, 2-Ethylhexyl mercaptoacetate and Linoleic acid ethyl ester have been confirmed to show antibacterial activities (Ajayi et al., 2011). This suggests that dietary supplementation of *Sauropus androgynous* essential oil can help to reduce the increasing cases of antimicrobial resistance and promote food safety (Ajayi et al., 2011). Paul et al. (2011), essential oil from *Sauropus androgynous* can defend against infection, disrupts cell membrane of pathogens and inhibited

Staphylococcus spp and *Salmonella spp* in antimicrobial assay. Preveen et al. (2011) reported that *Sauropus androgynous* oil is potent in

controlling bacterial growth in biofilms of *Streptococcus spp* and *Shigella spp*.

Compounds	R.T (min)	Molecular formula	Molecular weight (g/mol)	% Area
n-hexadecanoic acid	16.73	C ₁₆ H ₃₂ O ₂	256.424	10.45
Dipyrimadole	18.52	C ₂₄ H ₄₀ N ₈ O ₄	504.626	2.31
Octadec -9-enoic acid	18.90	C ₁₈ H ₃₄ O ₂	282.47	0.94
N-3-methylButyl acetamide	18.92	C ₇ H ₁₅ NO	129.20	1.46
Benzene,1,4-dichloro	21.13	C ₆ H ₄ Cl ₂	147.00	0.01
8-octadecenoic acid	22.46	C ₁₉ H ₃₆ O ₂	296.487	0.56
Humulene	22.78	C ₁₅ H ₂₄	204.35	0.18
Undecane, 2,10-dimethyl	22.98	C ₁₃ H ₂₈	184.36	0.46
Oxalic acid	24.06	C ₁₅ H ₂₈ O ₄	272.38	2.77
Nonane, 4-methyl	25.67	C ₁₀ H ₂₂	142.28	13.82
2-Ethylhexyl mercaptoacetate	25.75	C ₁₀ H ₂₀ O ₂ S	247.34	1.81
gamma. -Terpinene-3-Carene	26.31	C ₁₀ H ₁₆	136.23	2.28
Hexatriacontane	28.94	C ₃₆ H ₇₄	507.00	2.15
3-Isothiazolone	29.07	C ₃ H ₃ NOS	101.13	1.26
Nerolidol	29.35	C ₁₅ H ₂₆ O	222.37	0.14
10-Methylnonadecane	29.94	C ₂₀ H ₄₂	282.5	1.33
Cyclopropane	30.05	C ₃ H ₆	42.08	2.09
Ethyl palmitate	31.24	C ₁₈ H ₃₆ O ₂	284.5	0.71
2-methyl-Glutaric acid	31.88	C ₆ H ₁₀ O ₄	146.14	0.20
Linoleic acid ethyl ester	31.90	C ₁₈ H ₃₂ O ₂	308.5	0.38
9,17-Octadecadienol	33.25	C ₁₈ H ₃₂ O	264.4	0.05
4-Trifluoroacetoxytetradecane	33.57	C ₁₆ H ₂₉ F ₃ O ₂	424.4	0.57
Hexacosanoic acid	33.89	C ₂₆ H ₅₂ O ₂	396.7	1.10
Beta. -Famesene	34.21	C ₁₅ H ₂₄	204.35	1.59
Cyclopentaneundecanoic acid	34.57	C ₁₆ H ₃₀ O ₂	254.41	1.87
Pentadecane	37.21	C ₁₅ H ₃₂	212.41	0.01
Caryophyllene	37.67	C ₁₅ H ₂₄	204.351	13.92
Trans-Nerolidol	38.58	C ₁₅ H ₂₆ O	222.4	1.12
Methyl stearate	38.71	C ₁₉ H ₃₈ O ₂	298.5	1.07
Ethyl ricinoleate	39.12	C ₂₀ H ₃₈ O ₃	326.5	0.08
Squalene	39.65	C ₃₀ H ₅₀	410.7	19.63
Hexahydrofarnesyl acetone	40.19	C ₁₈ H ₃₆ O	268.5	1.24
Phytol	41.25	C ₂₂ H ₄₂ O ₂	296.5	8.71
Total				96.27
Number of compounds				33.00

Table 1: Bioactive compounds in *Sauropus androgynous* essential oil by GC-MS

Microorganisms	Ciproflaxacin	% Inhibition
<i>Staphylococcus aureus</i>	71	77.81
<i>Escherichia coli</i>	65	72.00
<i>Pseudomonas aeruginosa</i>	75	80.09
<i>Shigella flexineri</i>	68	73.73
<i>Salmonella typhi</i>	70	76.26

Table 2: Antimicrobial activity of *Sauropus androgynous* essential oil

Conclusion

It was concluded that *Sauropus androgynous* essential oil contains several unique bioactive compounds with therapeutic potentials and has been traditionally utilized for the treatment of cold, fever, inflammation, acute and chronic infections. *In vitro* studies have also shown that these compounds also have antimicrobial properties and was able to inhibit the activities of *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Shigella flexineri* and *Salmonella typhi*. Together they form a weapon against infection, inflammation and degeneration.

References

- Samad, A.P.A., U. Santoso, C.S. Chang and F.H. Nan, 2013. Identifying and analyzing bioactive components of *Sauropus*

androgynus ethanolic leaves extract. J. Fish. Soc. Taiwan, 40: 257-265

- Ajaiy, I.A., Ajibade, O. & Oderinde, R.A. (2011). Preliminary Phytochemical Analysis of some Plant Seeds. Research Journal of Chemical Sciences, 1 (3), 58-63.
- Ajiboye, A.A., Agboola, D.A., Fadimu, O.Y. & Afolabi, A.O. (2013). Antibacterial, Phytochemical and Proximate Analysis of *Prosopis africana* (LINN) Seed and Pod Extracts. FUTA Journal of Research in Sciences, 9 (1), 101-109.
- Lin, C.Q.; Lin, W.B.; Pan, W.D.; Lee, Y.J. Study on chemical constituents of the essential oil from the leaves of *Sauropus androgynus* L. Merr. J. Trop. Subtrop. Bot. 1999, 3, 255-256

5. Basker, A. Phytochemical Analysis and GC-MS Profiling in the Leaves of *Sauvopus androgynus* (L) MERR. *Int. J. Drug Dev. Res.* 2012, 4, 162–167.
6. Choi, J., Park, N., Hwang, S., Sohn, J., Kwak, I., Cho, K.K. & Choi, I. (2013). The antibacterial activity of various saturated and unsaturated fatty acids against several oral pathogens. *Journal of Environmental Biology*, 34, 673- 676.
7. Mamza, U.T., Sodipo, O.A. & Khan, I.Z. (2012). Gas Chromatography - Mass spectrometry (GC - MS) analysis of bioactive components of *Phyllanthus amarus* Leaves. *International Research Journal of Plant Science*, 3 (10), 208 - 215.
8. Awa, E.P., Ibrahim, S. & Ameh, D.A. (2012). GC/MS Analysis and Antimicrobial Activity of Dimethyl Ether fraction of Methanolic Extract from stem Bark of *Annona senegalensis* Pers. *International Journal of Pharmaceutical Sciences and Research*. 3 (11), 4213 – 4218.
9. Rajeswari Bose, Moorthy Saravana Kumar, Arumugam Manivel and Shanmugavadielv Chandra Mohan, 2018. Chemical Constituents of *Sauvopus androgynus* and evaluation of its antioxidant activity. *Res. J. Phytochem.*, 12: 7-13.
10. Singh, S.; Singh, D.R.; Salim, K.M.; Srivastava, A.; Singh, L.B.; Srivastava, R.C. Estimation of proximate composition, micronutrients and phytochemicals compounds in traditional vegetables from Andaman and Nicobar Islands. *Int. J. Food Sci.* 2011, 62, 765–773
11. Juxian, G.; Xian, Y. Studies on the Toxicology of *Sauvopus androgynus*, a Wild Vegetable in South China. *J. South China Agric. Univ.* 2005, 28, 10–14. Available online: <https://agris.fao.org/agris-search/search.do?recordID=CN2006002063> (accessed on 5 November 2022).
12. Purba, R.A.P.; Paengkoum, P. Exploring the Phytochemical Profiles and Antioxidant, Antidiabetic, and Antihemolytic Properties of *Sauvopus androgynus* Dried Leaf Extracts for Ruminant Health and Production. *Molecules* 2022, 27, 8580. <https://doi.org/10.3390/molecules27238580>
13. Andarwulan, N.; Kurniasih, D.; Apriady, R.A.; Rahmat, H.; Roto, A.V.; Bolling, B.W. Polyphenols, carotenoids, and ascorbic acid in underutilized medicinal vegetables. *J. Funct. Foods* 2012, 4, 339–347
14. Bao, H.; Sagues, W.J.; Wang, Y.; Peng, W.; Zhang, L.; Yang, S.; Xiao, D.; Tong, Z. Depolymerization of lignin into monophenolics by ferrous/persulfate reagent under mild conditions. *ChemSusChem* 2020, 13, 6582–6593.
15. Purba, R.A.P.; Paengkoum, S.; Paengkoum, P. Development of a simple high-performance liquid chromatography-based method to quantify synergistic compounds and their composition in dried leaf extracts of *Piper sarmentosum* Robx. *Separations* 2021, 8, 152.
16. Neamsuvan, O.; Ruangrit, T. A survey of herbal weeds that are used to treat gastrointestinal disorders from southern Thailand: Krabi and Songkhla provinces. *J. Ethnopharmacol.* 2017, 196, 84–93.
17. Petrus, A.J.A. *Sauvopus androgynus* (L.) Merrill-a potentially nutritive functional leafy-vegetable. *Asian J. Chem.* 2013, 25, 9425–9433.
18. Zhang, D.; Ivane, N.M.A.; Haruna, S.A.; Zekrumah, M.; Elysé, F.K.R.; Tahir, H.E.; Wang, G.; Wang, C.; Zou, X. Recent trends in the micro-encapsulation of plant-derived compounds and their specific application in meat as antioxidants and antimicrobials. *Meat Sci.* 2022, 191, 108842.
19. González-Romero, J.; Guerra-Hernández, E.J.; Rodríguez-Pérez, C. Chapter 19—Bioactive compounds from *Moringa oleifera* as promising protectors of in vivo inflammation and oxidative stress processes. In *Current Advances for Development of Functional Foods Modulating Inflammation and Oxidative Stress*; Hernández-Ledesma, B., Martínez-Villaluenga, C., Eds.; Academic Press: Cambridge, MA, USA, 2022; pp. 379–399.
20. Purba, R.A.P.; Paengkoum, S.; Yuangklang, C.; Paengkoum, P. Flavonoids and their aromatic derivatives in *Piper betle* powder promote in vitro methane mitigation in a variety of diets. *Cienc. Agropec.* 2020, 44, e012420
21. Paul Mariya, Beena Anto K. Antibacterial activity of *Sauvopus androgynus* (L.) Merr. *Internat. J. Plant Sci.* 2011; 6 (1): 189–192.
22. Praveen Kumar P, Kumaravel S, Lalitha C. Screening of antioxidant activity, total phenolics and GC-MS study of *Vitex negundo*. *African Journal of Biochemistry Research*. 2011. Vol. 4(7); pp.191 195
23. Carranza, M. G., Sevigny, M. B., Banerjee, D., and Fox-Cubley, L. (2015) 'Antibacterial Activity of Native California Medicinal Plant Extracts Isolated from *Rhamnus Californica* and *Umbellaria Californica*'. *Annals of Clinical Microbiology and Antimicrobials* 14 (1), 29
24. Caldeira, I., Pereira, R., Clímaco, M. C., Belchior, A., and De Sousa, R. B. (2004) 'Improved Method for Extraction of Aroma Compounds in Aged Brandies and Aqueous Alcoholic Wood Extracts using Ultrasound'. *Analytica Chimica Acta* 513 (1), 125-134
25. Chatrou, L. W., Pirie, M. D., Erkens, R. H., Couvreur, T. L., Neubig, K. M., Abbott, J. R., Mols, J. B., Maas, J. W., Saunders, R. M., and Chase, M. W. (2012) 'A New Subfamilial and Tribal Classification of the Pantropical Flowering Plant Family Annonaceae Informed by Molecular Phylogenetics'. *Botanical Journal of the Linnean Society* 169 (1), 5–40
26. Gao, S., Lewis, G. D., Ashokkumar, M., and Hemar, Y. (2014) 'Inactivation of Microorganisms by Low-Frequency High-Power Ultrasound: 1. Effect of Growth Phase and Capsule Properties of the Bacteria'. *Ultrasonics Sonochemistry* 21 (1), 446-453
27. John AO. *Clerodendron splendens* leaf extract supplementation in weaner rabbits: impact on growth performance, haematology and intestinal microbial population. *Cerrado: Agricultural and Biological Research*, 2024b;1(1): 21-31.
28. John AO. Effect of coconut shell extract on the growth performance and some haemato-biochemical parameters of broiler chicken. *Brazilian Journal of Science*, 2024c; 3(6): 82-95.
29. John AO. Impact of dietary supplementation of *Rhamnus prinoides* leaf extract on the growth performance, nutrient retention and intestinal microbial count of Japanese quails. *Brazilian Journal of Science*, 2024a; 3(5): 40-50.
30. John AO. Effect of performance, serum biochemistry and hematological components of feeding Japanese quails phytopreparations comprising *Megaphrynum macrostachyum* leaves. *Brazilian Journal of Science*, 2024d; 3(5): 51-64.
31. Singh S, Alagbe OJ, Liu X, Sharma R, Kumar A. Comparative analysis of ethanolic *Juniperus thurifera* leaf, stem bark and root extract using gas chromatography and mass spectrometry. *International Journal of Agriculture and Animal Prod.* 2022; 2(6): 18-27.
32. Singh AS, Alagbe JO, Sharma S, Oluwafemi RA, Agubosi OCP. Effect of dietary supplementation of melon (*Citrullus lanatus*) seed oil on the growth performance and antioxidant status of growing rabbits. *Journal of Multidimensional Research and Rev.* 2021; 2(1): 78-95.
33. Hernandez M, Alagbe JO. (2025). Influence of *Abrus precatorius* crude oil supplementation on the growth performance, Nutrient digestibility, Ruminal fermentation and

Microbial population of Malabari Bucks. International Journal of Global Sustainable Res. 2025a; 3(7): 527-538

34. Hernandez M, Alagbe JO. (2025). Influence of *Odontonema strictum* oil on the growth performance and Ruminal fermentation of Barbary bucks. Research in: Agriculture and Veterinary Sci. 2025b; 9(2):1-10.

35. Muritala, DS, Daniel Shittu, Alagbe, JO, Ojebiyi OO, Ojediran TK, Rafiu TA. Growth performance and haematological and serum biochemical parameters of broiler chickens given varied concentrations of *Polyalthia longifolia* leaf extract in place of conventional antibiotics. *Animal Science and Genetics* 2022; 18(2): 57-71.

36. Ojediran TK, Emiola IA, Durojaye, V, Alagbe, JO. Analysis of *Kigelia africana* fruit's powder antioxidant and phytochemical properties. *Brazilian Journal of Science*, 2024; 3(7): 38-49.

37. Ojediran TK, Emiola IA, Durojaye V, John O.A. Proximate, vitamin and GC-MS profiling of *Kigella africana* powder. *Cerrado: Agricultural and Biological Research*, 2024; 1(1): 13-20.

38. Shittu MD, Alagbe JO, Alaba O, Okanlawon EO, Adelakun FA, Emmanuel EO and Adejumo DO. Effect of ginger, garlic and Negro pepper on the gut microbes, gut histomorphometry and pathological assessment of selected organs of broiler chickens. *Association of Deans of Agriculture in Nigerian Universities*, 2024; 5: 105-121.

39. Oluwafemi RA, Lawal A.O, Adelowo SA, Alagbe JO. Effects of dietary inclusion of ginger (*Zingiber officinale*) and garlic (*Allium sativum*) oil on carcass characteristics and sensory evaluation of broiler chicken. *Texas Journal of Multidisciplinary Stud*. 2021; 2(11): 180-188.

40. Alagbe JO, Shittu MD, Tanimomo, Babatunde K. Influence of *Anogeissusleio carpus* stem bark on the fatty acid composition in meat of broiler chickens. *European Journal of Life Safety and Stability* 2022; 14(22): 13-22.

41. Alagbe JO. Use of medicinal plants as a panacea to poultry production and food security: A review. *Gospodarka I Innowacje* 22(2022): 1-12.

42. Omokore EO, Alagbe, JO. Efficacy of dried *Phyllanthus amarus* leaf meal as an herbal feed additive on the growth performance, haematology and serum biochemistry of growing rabbits. *International Journal of Academic Research and Development*.2019; 4(3): 97-104.

43. Musa B, Alagbe JO, Adegbite MB, Omokore EA. Growth performance, caeca microbial population and immune response of broiler chicks fed aqueous extract of *Balanites aegyptiaca* and *Alchornea cordifolia* stem bark mixture. *United Journal for Research and Technology*, 2020; 2(2):13-21.

44. Adewale AO, Alagbe JO, Adeoye, AO. Dietary Supplementation of *Rauvolfia Vomitoria* Root Extract as A Phytonic Feed Additive in Growing Rabbit Diets: Haematology and serum biochemical indices. *International Journal of Orange Technologies*, 2021; 3(3): 1-12.

45. Alagbe OJ. Novel phytonic's impact on weaned pigs growth performance, haematology and serum biochemical indicators. *Black Sea Journal of Agriculture*, 2024; 7(2): 82-89.

46. Shittu MD, Alagbe JO. (2020) Phyto-nutritional profiles of broom weed (*Sida acuta*) leaf extract. *International Journal of Integrated Edu* 2020; 3(11): 119-124

47. Alagbe JO. (2022). Gas chromatography and mass spectroscopy of *Juniperus phoenicea* stem bark extract and its influence on the haemato-biochemical values of growing rabbits. *British Scientific Period*. 2022; 1(1): 18-33.

48. Alagbe JO, Zubairu H, Adedeji OM, Bamigboye S, Dora A. Influence of *Juniperus thurifera* root extract on the nutrient digestibility and caecal microbial count of growing rabbits. *Web of Synergy: International Interdisciplinary Res Jou*. 2022; 1(1): 5-17.

49. NIST. Chemistry Web Book. Data from NIST Standard Reference Database 69, 2011.

50. Sharma, R.R. (2012). Preliminary phytochemical screening of some indigenous medicinal plant leaves extract in regulation of antidiabetic activity. *Science Research Reporter*, 2 (3), 307 310.

51. Jing-Chung, C., Jang-Ying, Y., Pei-Chun, C. & Cheng-Kuang, H. (2007). Phenolic content and DPPH radical scavenging activity of Yam containing Surimi gels influenced by salt and heating. *Asian Journal of Health and Information Sciences*, 2, 1-4, 1-11.

52. Lima, A.L., Parial, R., Das, M. & Das, A.K. (2010). Phytochemical and Pharmacological studies of ethanolic extract from the leaf of mangrove plant *Phoenix paludosa* Roxb. *Malaysian Journal of Pharmaceutical Sciences*, 8 (2), 59-69.

53. Manilal, A., Sujith, S., Sabarathnam, B., Kiran, G.S., Selvin, J., Shakir, C. & Lipton, A. (2011). Biological activity of the red algae *Laurencia brandenii*. *Acta Botanica Croatica*, 70 (1), 81-90.

54. Namkeleja, H.S., Tarimo, N.T.C. & Ndakidemi, P.A. (2014). Allelopathic Effects of *Argemone Mexicana* to Growth of Native Plant Species. *American Journal of Plant Sciences*, 5 (9), 1336 1344.

55. Tiwari, P., Kumar., B., Kaur, M., Kaur, G. & Kaur, H. (2011). Phytochemical screening and Extraction: A Review: *International pharmaceutica sciencia*, 1 (1), 98-106.

56. Sinha, M.K. & Munshi, J.D. (2010). Toxicology of biocidal plants. *Delhi, Mittal Publications, India*; pp53.

57. Zubairu, H., Mubarak, A. A and Alagbe, J.O. (2025). Effect of partial replacement of Soymeal with *Sphenocentrum jollyanum* seed meal on the growth performance and haematological constituents of broiler chickens. *International Journal of Integrative Research*, 3(10): 741-752.

58. Alagbe, J.O. (2025). Assessing the bioactive compounds of *Thaumatomoccus daniellii* leaf extract by GC-MS. *International Journal of Global Sustainable Research*, 3(9): 617-626.

59. Agubosi, O.C.P., Alexander, James and Alagbe, J.O. (2022). Influence of dietary inclusion of Sunflower (*Helianthus annus*) oil on growth performance and oxidative status of broiler chicks. *Central Asian Journal of Medical and Natural Sciences* 2(7): 187-195.

60. Agubosi, O.C.P., Soliu, M.B and Alagbe, J.O. (2022). Effect of dietary inclusion levels of *Moringa oleifera* oil on the growth performance and nutrient retention of broiler starter chicks. *Central Asian Journal of Theoretical and Applied Sciences* 3(3): 30-39.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here:

[Submit Manuscript](#)

DOI:[10.31579/2637-8914/346](https://doi.org/10.31579/2637-8914/346)

Ready to submit your research? Choose Auctores and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At Auctores, research is always in progress.

Learn more <https://auctoresonline.org/journals/nutrition-and-food-processing>